

Synthesis and Characterization of 2-[1h- Benzimidazole- 2yl-Sulfanyl]-N-[(E)-Furan-2-Yl- Methylidene] Aceto Hydrazide

*¹Ramesh Dhani

¹Department of Pharmaceutical Chemistry, Oil Technological Research Institute, Jawaharlal Nehru Technological University Anantapur, Andhra Pradesh, India

Manuscript No: IJPRS/V1/I4/00185, Received On: 11/10/2012, Accepted On: 17/10/2012

ABSTRACT

Heterocyclic chemistry comprises at least half of all organic chemistry research worldwide. In particular, heterocyclic structures form the basis of many pharmaceutical, agrochemical and veterinary products. The benzimidazole contains a phenyl ring fused to an imidazole ring, as indicated in the structure of benzimidazole. Diversity of biological response profile has attracted considerable interest of several researchers across the globe to explore this skeleton for its assorted therapeutic significance. By using different synthetic methods new benzimidazole derivatives were synthesized and further Melting points were determined by using Precision melting point apparatus in open capillaries and are uncorrected. The purity of the compounds was checked by TLC on silica gel G plates using n-Hexane, ethyl acetate (1:3) and methanol: chloroform (1:9) solvent system. The synthesized benzimidazole derivatives were characterized by IR, ¹H NMR spectral analysis. Benzimidazole is a lead nucleus for future developments to get effective compounds.

KEYWORDS

Benzimidazole, Furan, Acetohydrazide, Imidazole, n-Hexane, Ethyl acetate, Chloroform, Methanol.

INTRODUCTION

Many important biochemical compounds and drugs of natural origin contain heterocyclic ring structures. Among these e.g. Carbohydrates, essential amino acids, vitamins, alkaloids, glycosides etc. the presence of heterocyclic structures in such diverse types of compounds is strongly indicative of the diverse types of the pharmacological activity and recognition of this is reflected in efforts to find useful synthetic drugs. Medicinal chemistry, according to Berger, “tried to be based on the ever increasing hope that biochemical rationale for drug discovery may be found”.

Pharmaceutical chemistry is a science that makes use of general laws of chemistry to study drugs, i.e. their preparation, chemical nature, composition, structure, influence on an organism and studies of the physical and chemical properties of drugs, the methods of quality control and the conditions of their storage. Heterocyclic chemistry is the branch of chemistry dealing with the synthesis, properties, and applications of heterocyclics.

Heterocyclic chemistry comprises at least half of all organic chemistry research worldwide. In particular, heterocyclic structures form the basis of many pharmaceutical, agrochemical and veterinary products. The benzimidazole contains a phenyl ring fused to an imidazole ring, as indicated in the structure of benzimidazole. The important group of substances has found practical application in a number of fields. Recently in benzimidazole chemistry has been

***Address for Correspondence:**

Ramesh Dhani

Department of Pharmaceutical Chemistry,
Oil Technological Research Institute,
Jawaharlal Nehru Technological University Anantapur,
Andhra Pradesh, India.

E-Mail Id: dhani.ramesh1@gmail.com

revived somewhat by the discovery that the 5, 6-dimethyl benzimidazole moiety is a part of the chemical structure of vitamin B12.

MATERIALS AND METHODS

1. Synthesis of 2-mercapto benzimidazole

A mixture of 10.8gm (0.1mol) of o-phenylenediamine, 5.65 gm (0.1mol) of potassium hydroxide and 7.67 gm (0.1mole, 6.19ml) of carbon disulfide, 100ml of 95% ethanol and 15 ml of water was taken in a 500ml round bottom flask heated under reflux for three hours. Then 1-1.5 gm of charcoal was added cautiously and the mixture is further heated at the reflux for 10 minutes, the charcoal is removed by filtration. The filtrate is heated to 60-70°C, 100ml of warm water is added, and acidified with dilute acetic acid with good stirring. The product separated as glistening white crystals, and the mixture is placed in a refrigerator for three hours to complete the crystallization. The product is collected on a Buckner funnel and dried over night at 40°C. The dried product is recrystallised by ethanol the yield is 8.5gm (73%) melting point is 300-305°C.




Table 1: Physicochemical analysis

S.no	2-mercapto benzimidazole	
1.	Mol. Formula	C ₇ H ₆ N ₂ S
2.	Melting Point	300-305°C
3.	% Yield	73%
4.	Solvent system used	hexane: ethyl acetate (1:3)

2. Synthesis of ethyl (1H-benzimidazol-2-yl-sulfanyl) acetate

A stirred mixture containing 4.5gm of (0.03mole) of 2-mercaptopbenzimidazole, 60ml of ethanol and 1.68gm of (0.03mole) potassium hydroxide was added and heated at 78-80°C for 10-minutes. Then ethyl chloro acetate (3.66ml, 0.03mol) was added in one portion, an exothermic reaction set in causing a temperature rise from 30-40°C. After stirring at 25-30°C for 18-hours, the reaction mixture was added to 100gm of ice-water and stirred for 30-minutes at 0-10°C. The precipitate was collected by filtration washed with water until free of chloride and air dried at 50°C and recrystallised by water the yield is 6 gm (62.25%). melting point is 105°C.




Table 2: Physicochemical analysis

S.no	Ethyl (1H-benzimidazol-2-yl-sulfanyl) acetate	
1.	Mol. Formula	C ₁₁ H ₁₂ O ₂ N ₂ S
2.	Melting Point	105°C
3.	% Yield	62.25 %
4.	Solvent system used	hexane: ethyl acetate (1:3)

3. Synthesis of 2-(1H-benzimidazol-2-ylsulfanyl) acetohydrazide

The mixture of 2-carboxy ethyl thio 1H-benzimidazole 4gm (0.004mole) and hydrazine

hydrate 6ml (0.01mole) are mixed well in a RBF and heated on water bath for 10 min. then dissolved in 60 ml ethanol, the reaction mixture is heated with reflux the reaction mixture is heated with reflux condenser for six hours, cooled to room temperature and the reaction mixture was added to 100gm of ice-water, and kept aside for the crystallization. The colorless crystals are collected by filtration, and recrystallized from water. Melting point is 180-185°C; the yield is 60-70%.





Table 3: Physicochemical analysis

S.no	2-(1H-benzimidazol-2-yl-sulfanyl)acetohydrazide	
1.	Mol. Formula	$\text{C}_8\text{H}_{11}\text{ON}_4\text{S}$
2.	Melting Point	180-185°C
3.	% Yield	60-70%
4.	Solvent system used	hexane: ethyl acetate (1:3)

4. General procedure for the preparation of schiff bases compound



A equimolar solution of carboxyl hydrazide(0.009 mol, 2gm) is dissolved in 10ml of ethanol and to this solution substituted aldehydes in equi molar qty (0.009mol, 0.917) is added with 4-6 drops of glacial acetic acid was added, this reaction mixture is kept under reflux for 8 hours. After cooling to room temperature was added to cold water. Compound gets separated as solid filtered, dried and recrystallized with chloroform.




Table 4: Physicochemical analysis

S.no	2-[1H- benzimidazol- 2yl-sulfanyl]-N-[(E)-furan-2-yl-methylidene] acetohydrazide	
1.	Mol. Formula	$\text{C}_{14}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$
2.	Melting Point	230-235°C
3.	% Yield	65.5%
4.	Mol. Weight	300

RESULTS AND DISCUSSION

The present study explains the synthesis and characterization of benzimidazol derivative i.e., 2-[1H- benzimidazol- 2yl-sulfanyl]-N-[(E)-furan-2-yl-methylidene]acetohydrazide. At present studies find the structural-activity relationship (SAR) and to optimize the structure. The synthesized benzimidazol derivative characterized by IR, ^1H NMR spectral data analysis. The purity of the synthesized benzimidazol derivative was checked by (TLC) thin layer chromatography and R_f value was recorded.

Table 5: IR spectral data

Functional group assigned	Group frequency in Wave number (cm ⁻¹)
(-NH-)	3333,1341
(>C=O)	1676
(-C=N-)	1618
C-S-C	681
aromatic ring	3074




Figure 1: IR Spectra of 2-[1H- benzimidazole- 2yl-sulfanyl]-N-[(E)-furan-2-yl-methylidene] acetohydrazide




Figure 2: ¹H NMR Spectra of 2-[1H benzimidazole- 2yl-sulfanyl]-N-[(E)-furan-2-yl-methylidene] acetohydrazide

Table 6: ¹H NMR spectral data

S. no	Value (δ)	Nature of segment	Type
1	11.9	Singlet	NH of Benz
2	9.6	Singlet	1 H, NH-N
3	8.0	Singlet	1H, N=CH
4	6.7-7.7	Multiplet	7H, Ar H
5	4.7	Singlet	2H, S-CH ₂

CONCLUSION

By this study concluded that to find the structure-activity relationship (SAR) and to optimize the structure of the synthesized new benzimidazol derivative i.e., 2-[1H-benzimidazole-2yl-sulfanyl]-N-[(E)-furan-2-yl-methylidene] acetohydrazide. The compound was characterized by IR, ¹H NMR spectral data, the purity of the compound was checked by TLC and it produces good yield. The compound was confirmed by physicochemical and spectral data analysis.

ACKNOWLEDGEMENT

I am indebted to my parents and my sister for their inspiration and encouragement given to me during this work with deep appreciation for their determination and enthusiasm at each and every front of my life to transform my dreams into reality. I am very thankful and prevail age to my deep sense of gratitude to Prof. Dr. Bishwaranjan Behra, M.Pharm., Ph.D.

REFERENCES

1. Kubo K et. al., Synthesis of 2-[[4-fluoroalkoxy -2-pyridyl] methyl] sulfinyl] -1H-benzimidazole as anti-ulcer agents. ChemPharma Bull. 1900, 38, 2853-2858.
2. Chakraborty, P. In A Text Book of Microbiology, Ed 1, Orient Longman Publisher, 2000, pp 3-20, 73-78, 378-382.
3. Dejongh DC and Thomson ML. Pyrolysis and mass spectra of the 2-thiones of benzothiazole, benzimidazole and benzoxazole. Journal of Organic Chemistry. 1973, 38 (7), 1356-1361.

4. Padmaja J et. al., Reaction of 2-(1-amino-2-naphthyl) benzimidazole with aldehydes. Indian Journal of Chemistry. 1988, 27, 418-420.
5. John B Wright et. al. The chemistry of benzimidazoles. Chemical Reviews, 1951, 3(3), 397-541.
6. Day AR. Electronic mechanism of organic synthesis. 1951, 2, 242-243.
7. Preston PN. Synthesis Reactions and Spectroscopic properties of benzimidazoles. Chemical Reviews. 1974, 15, 279-314.
8. Wagner EC and Millett WH. Synthesis of Benzimidazole. Chemistry of organic Synthesis. 1943, 2, 65.
9. Shriner RL and Robert W Upson. Synthesis of bis-benzimidazoles from diabasic acids. Journal of Organic Chemistry. 1941, 63(8), 2277.
10. Walther R Von and Kessler A. Journal of Prakt Chemistry. 1906, 174, 245.
11. Goodman Gilman, Alfred and McGraw-Hill, The Pharmacological Basis of Therapeutics, ed10, McGraw-Hill Medical Publishing Division, 2001, 1284.
12. Fromling A, Recent Trends in the Discovery, Development and Evaluation of Antifungal Agents, Ed 7, Prous Barcelona, 1987, 12-25.

